



Giannakopoulos, A. and Valiakos, G. and Papaspyropoulos, K. and Dougas, G. and Korou, L. M. and Tasioudi, K. E. and Fthenakis, G. C. and Hutchings, M. R. and Kaimaras, D. and Tsokana, C. N. and Iliadou, P. and Spyrou, V. and Tzani, M. and Birtsas, P. and Kostoglou, P. and Sokos, C. and Doudounakis, S. and Yon, L. and Hannant, Duncan and Artois, M. and Tsiodras, S. and Hadjichristodoulou, C. and Billinis, C. (2016) Rabies outbreak in Greece during 2012–2014: use of Geographical Information System for analysis, risk assessment and control. *Epidemiology and Infection*, 144 (14). pp. 3068-3079. ISSN 1469-4409

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/37950/1/Rabies_Epid-Inf%20%28002%29.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk



Rabies outbreak in Greece during 2012-2014: use of Geographical-Information-System for analysis, risk assessment and control

Journal:	<i>Epidemiology and Infection</i>
Manuscript ID	HYG-OM-7137-Feb-16.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	17-May-2016
Complete List of Authors:	<p>Giannakopoulos, Alexios; University of Thessaly, Faculty of Veterinary Medicine</p> <p>Valiakos, George ; University of Thessaly, Faculty of Veterinary Medicine</p> <p>Papaspyropoulos, Konstantinos ; Hunting Federation of Macedonia and Thrace, Research Division</p> <p>Douglas, Georgios ; Hellenic Centre for Disease Control and Prevention</p> <p>Korou, Laskarina-Maria ; Animal Health Directorate, Ministry of Rural Development and Food, Department of Zoonoses</p> <p>Tasioudi, Konstantia ; Athens Veterinary Centre, Ministry of Rural Development and Food,, Department of Molecular Diagnostics, FMD, Virological, Rickettsial and Exotic diseases</p> <p>Fthenakis, George; School of Veterinary Medicine, ;</p> <p>Hutchings, Michael; SRUC, Disease Systems</p> <p>Kaimaras, Dimitrios ; Ministry of Rural Development and Food, Directorate of Technical Studies Structures and Topography</p> <p>Tsokana, Constantina; University of Thessaly, Faculty of Veterinary Medicine</p> <p>Iliadou, Peristera ; Athens Veterinary Centre, Ministry of Rural Development and Food,, Department of Molecular Diagnostics, FMD, Virological, Rickettsial and Exotic diseases,</p> <p>Spyrou, Vassiliki ; Technological Education Institute of Thessaly, Department of Animal Production</p> <p>Tzani, Myrsini ; Animal Health Directorate, Ministry of Rural Development and Food, Department of Zoonoses</p> <p>Birtsas, Periklis ; Technological Institute of Thessaly, Department of Forestry and Natural Environment Administration</p> <p>Kostoglou, Petroula ; Animal Health Directorate, Ministry of Rural Development and Food, Department of Zoonoses</p> <p>Sokos, Christos ; University of Thessaly, Faculty of Veterinary Medicine</p> <p>Doudounakis, Spyros ; Hellenic Ministry of Rural Development and Food, Animal Welfare Identification and Veterinary Drugs and Residues Department</p> <p>Yon, Lisa; University of Nottingham, School of Veterinary Medicine & Science</p> <p>Hannant, Duncan; University of Nottingham, Faculty of Medicine and Health Sciences</p> <p>Artois, Marc ; University of Lyon, VetAgro Sup</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	Tsiodras, Sotirios; University of Athens Medical School, Athens, Greece., 4th University Dept of Internal Medicine HADJICHRISTODOULOU, CHRISTOS; University of Thessaly, Medicine Billinis, Charalambos; School of Veterinary Medicine, Laboratory of Microbiology and Parasitology
Keyword:	fox, geographical-information-system, Greece, rabies

SCHOLARONE™
Manuscripts

For Review Only

Rabies outbreak in Greece during 2012-2014: use of Geographical-Information-System for analysis, risk assessment and control

A. GIANNAKOPOULOS^{1†}, G. VALIAKOS^{1†}, K. PAPASPYROPOULOS², G. DOUGAS³, L.M. KOROU⁴, K. E. TASIOUDI⁵, G. C. FTHENAKIS¹, M. R. HUTCHINGS⁶, D. KAIMARAS⁷, C. N. TSOKANA¹, P. ILIADOU⁵, V. SPYROU⁸, M. TZANI⁴, P. BIRTSAS⁹, P. KOSTOGLOU⁴, C. SOKOS¹, S. DOUDOUNAKIS¹⁰, L. YON¹¹, D. HANNANT¹¹, M. ARTOIS¹², S. TSIODRAS¹³, C. HADJICHRISTODOULOU^{3,14}, C. BILLINIS^{1*}

¹ Faculty of Veterinary Medicine, University of Thessaly, Karditsa, Greece

² Research Division, Hunting Federation of Macedonia and Thrace, Thessaloniki, Greece

³ Hellenic Centre for Disease Control and Prevention, Athens, Greece

⁴ Department of Zoonoses, Animal Health Directorate, Ministry of Rural Development and Food, Athens, Greece

⁵ Department of Molecular Diagnostics, FMD, Virological, Rickettsial and Exotic diseases, Athens Veterinary Centre, Ministry of Rural Development and Food, Athens, Greece

⁶ Disease Systems, Scotland's Rural College, Edinburgh, United Kingdom

⁷ Directorate of Technical Studies, Structures and Topography, Ministry of Rural Development and Food, Athens, Greece

⁸ Department of Animal Production, Technological Education Institute of Thessaly, Larissa, Greece

⁹ Department of Forestry and Natural Environment Management, Technological Institute of Thessaly, Karditsa, Greece

¹⁰ Animal Welfare Identification and Veterinary Drugs and Residues Department, Animal Health Directorate, Ministry of Rural Development and Food, Athens, Greece

¹¹ School of Veterinary Medicine & Science, University of Nottingham, Nottingham, United Kingdom

¹² University of Lyon, VetAgro Sup, France

¹³ Fourth Academic Department of Internal Medicine, University of Athens Medical School, Athens, Greece

¹⁴ Faculty of Medicine, University of Thessaly, Larissa, Greece

*Author for correspondence: Prof. C. Billinis, Department of Microbiology and Parasitology, Faculty of Veterinary Medicine, University of Thessaly, 224 str. Trikalon, 43100, Karditsa, Greece

E-mail: billinis@vet.uth.gr

† These authors have contributed equally to the work and their names are listed alphabetically.

Short running head: Investigation of rabies outbreak in Greece using GIS

SUMMARY

Objectives of this work were (i) geographic analysis of the 2012-2014 outbreak of rabies in Greece using GIS and (ii) comparative analysis of animal cases with data of potential human exposure to rabies together with environmental data, in order to provide information for risk assessment, effective monitoring and control. Most animal cases (40/48) involved red foxes, while domestic animals were also diagnosed with rabies. Overall, 80% of the cases were diagnosed in central northern Greece; 75% of the cases were diagnosed in low altitudes (<343.5 m), within a distance of 1 km from human settlements. Median distance from livestock farms was 201.25 m. Most people potentially exposed to rabies (889/1060) were presented with dog bite injuries. Maximum entropy analysis revealed that distance from farms contributed with the highest percent to define environmental niche profiles for rabid foxes. Oral vaccination programs were implemented in 24 administrative units of the country during 2013 and 2014, covering a total surface of approx. 60,000 km². Rabies reoccurrence in Greece emphasizes the need for ongoing surveillance in cross-border areas and in areas with intense human activity.

Key words: fox, geographical-information-system, Greece, rabies

INTRODUCTION

Rabies, an acute progressive disease of the central nervous system caused by neurotrophic viruses of the genus *Lyssavirus*, is almost always fatal in humans without timely medical care (e.g., post-exposure prophylaxis); every year, 55000 people around the world die of the disease. In Europe, rabies is found mainly in wild animals [1]. In Eastern Europe, at least four different variants of the causal virus have been described, which belong to the groups CE (Central Europe), EE (Eastern Europe), NEE (North Eastern Europe) or SF (Serbian fox) [2, 3].

Greece borders with Albania, Former Yugoslav Republic of Macedonia (FYROM), Bulgaria and Turkey. In the last three countries, rabies has been reported in wild and domestic animals and wildlife vaccination campaigns using oral live attenuated vaccines are being implemented most recently: in Bulgaria since 2009, in FYROM since 2011 and in Turkey since 2008 [4-6]. Despite successful oral vaccination campaigns of wildlife

targeting to elimination of rabies in large parts of Europe [1], the disease still occurs in the continent and, in 2012, 4884 cases were diagnosed in animals.

Prior to 1950, rabies was endemic in Greece, although there were no official records regarding frequency of the disease. During the years 1953 and 1954, 995 and 1135, respectively, animal rabies cases were diagnosed, as well as six and four, respectively, rabies-related human deaths. In total, during the period 1951 to 1980, 11,472 animal rabies and 53 human rabies cases had been diagnosed. The last case in humans was diagnosed in 1970. Eradication of the disease has been attributed to (i) widespread vaccination of 'ownerless free-roaming' dogs, purposefully implemented, (ii) widespread, free of charge vaccination of domestic dogs and (iii) increased public awareness. Thereafter, only one single case of rabies was diagnosed in a hunting dog in the Evros area, near the border with Turkey, in 1987 [7, 8] (Fig. 1).

After 1987, Greece had been considered as a rabies-free country, although there were outbreaks of the disease in neighbouring countries [8], which might have contributed to re-introduction of the disease into Greece. In November 2011, a new case of rabies was detected in FYROM, 300 m from the Greek borders, in a dead red fox. Then, in October 2012, an outbreak of rabies was recorded within the Greek territory. Since then, many cases of the disease have been diagnosed in animals, mainly in red foxes (*Vulpes vulpes*), in the country. Phylogenetic analysis of virus isolates showed that these belong to EE group [7].

Red foxes are adaptable omnivorous carnivores distributed across all continents in the northern hemisphere. They have an important ecological and socio-economic role as a game species, as well as a key wildlife host of rabies virus [9]. Despite absence of a population estimate in Greece, red foxes are considered to have widespread distribution and high abundance in the country [10]. The adaptable and opportunistic nature of the species has enabled these animals to inhabit suburban and urban areas of Greece. Their close proximity with human populations and the resulting possible close contact of red foxes with ownerless free-roaming dogs or cats, for which there is limited management throughout Greece, support the hypothesis that red foxes were key contributors to rabies risk for public health.

Objectives of this work were (i) geographic analysis of the 2012-2014 outbreak of animal rabies in Greece using GIS and (ii) comparative analysis of animal cases with data of potential human exposure to rabies together with environmental data, in order to provide information for risk assessment, effective monitoring and control.

METHODS

Study area

Greece borders with Albania to the northwest, FYROM and Bulgaria to the north and Turkey to the northeast. Total surface of the country is 131990 km². The study area includes the local administration units, where cases of rabies have been diagnosed in animals, as well as administration units adjacent to those. Total surface of that area is 65167 km².

Greece is characterized by (i) presence of large mountain areas (approx. 80% of total surface of the country), (ii) extensive coastline (approx. 15000 km), (iii) many island complexes in the Aegean and the Ionian seas and (iv) large climatic diversity (29 climatic zones according to the Thorn Waite classification). In general, climatic conditions in the country are typically Mediterranean: summer is hot and dry and winter is usually mild. Rainfall occurs mostly in autumn and winter.

Passive surveillance of the disease and cases of rabies in animals

An already enforced national program for passive surveillance of rabies, carried out according to Directive 99/2003/EC, which provided collection and examination of wild and domestic animals found dead or suspected of having rabies, was enhanced in 16 prefectures along the northern and eastern land borders of Greece, after November 2011, when a fox rabies case was officially reported by FYROM near the Northern Greek borders. After laboratory confirmation of a rabid fox in October 2012 in Greece, collection of samples under the passive surveillance program was extended in 2013 throughout the country, with the objective to achieve a more efficient surveillance of the disease and to obtain more accurate results on circulation of the virus in the country.

Notification of suspect rabies cases according to the Greek legislation on rabies control makes mandatory by any person working with animals (farmers, private veterinarians, staff of local authorities responsible for control of ownerless free-roaming dogs and cats etc.) to immediately report any case of suspicion of rabies in animals. In addition, a public awareness campaign, conducted by the Animal Health Directorate of the Ministry of Rural Development and Food, targeted the general public, aimed to increase awareness and necessity for notification of all suspect cases. Further, game wardens and forestry officers were involved in collection of dead wild animals or animals suspect for rabies and their delivery to the competent regional veterinary authorities for subsequent appropriate submission to the national reference laboratory for animal rabies. The national rabies control and eradication program in animals, as well as the passive surveillance program for the disease were coordinated by the Animal Health Directorate of the Ministry of Rural Development and Food (i.e., the official veterinary service of the country).

Details of animal rabies cases (October 2012-June 2014) and information regarding organisation and implementation of the wildlife oral vaccination campaigns have been provided by the Animal Health Directorate. Laboratory diagnosis of rabies in animals is undertaken at the Athens Veterinary Centre of the Ministry of Rural Development and Food, where the Greek national reference laboratory for rabies in animals is located. Geographical coordinates of the locations, where cases of rabies were recorded, were obtained from official game wardens.

Cases of human potential exposure to rabies

Shortly after the identification of the first rabid animal, an algorithm including recommendations for management of human cases was provided to physicians. Criteria included geographic location (high, medium, low risk area), category of exposure, animal species involved, availability of the animal for evaluation in a defined time margin, presentation of the animal during incident and provoked or justified attack. Simultaneously, a notification system was established for data collection regarding the human cases potentially exposed to rabies. Details of people who were potentially exposed to rabies in Greece from October 2012 to June 2014, have been provided by the Hellenic Centre for Disease Control & Prevention (HCDCP), a service of

the Ministry of Health. Exposure might have arisen from contact with (i) animals (wild, ownerless free roaming or domestic) suspected to be rabies-infected or (ii) rabies vaccine baits.

Relevant data that were collected, included type of exposure, classification of exposure, animal species suspected to be rabies-infected involved (wildlife or domestic animal), location of the incident and type of anti-rabies treatment. For classification of exposure, the guidelines of the World Health Organisation were used; class III exposure involved mostly transdermal bites, single or multiple, class II exposure involved mostly minor scratches or abrasions with no bleeding and class I exposure involved touching or licking on intact skin. In relation to location of the incident, risk assessment of rabies circulation in the geographical area was also taken into account; in high risk areas (i.e., areas within a 50 km radius from the location where a case of rabies was diagnosed), treatment administration was always recommended; in medium risk areas (i.e., all other areas of administrative units in Northern Greece that were not classified as high risk), treatment administration was recommended only when, according to the description of the exposed person, the animal was considered suspect of rabies (non-provoked attack and/or signs [as described by the person affected] compatible with rabies); finally, in low risk areas (i.e., all areas that were not classified as high or medium risk), treatment administration was recommended only when, according to the description of the exposed person and with additional evaluation by a veterinarian the animal was suspect of rabies (non-provoked attack and/or signs compatible with rabies).

Oral anti-rabies vaccination areas

The area where oral anti-rabies vaccination programmes had been applied, was determined by the Animal Health Directorate, in collaboration with the Directorate of Technical Studies, Structures and Topography, also a service of the Ministry of Rural Development and Food, in accordance with EU guidelines and recommendations. The selected areas included (i) a 50 km ‘buffer zone’ around the location of each laboratory confirmed rabies case within the Greek territory, (ii) each local administrative unit where cases of animal rabies had been diagnosed and (iii) all adjacent to above administrative units.

According to the national Greek oral rabies vaccination (ORV) program for immunization of wildlife, which was issued before initiation of each ORV campaign and was approved by the relevant EU services, all urban and

suburban areas had been excluded of target areas of vaccination. Vaccine-baits were aerielly distributed by fixed-wing aircrafts. The method did not allow operations in areas with high density inhabitation [11]. An additional parametre contributing to the choice of the vaccination area, was the potential of increased human exposures to the vaccine used. The vaccine selected for the first oral vaccination campaign in Greece was the SAG2 vaccine (Rabigen® SAG2; Virbac, Carros, France), a live attenuated rabies virus (strain SAG2) vaccine. As the vaccine contained a live attenuated strain of the virus and there was lack of studies regarding safety of the vaccine to humans after direct contact, exposure to the vaccine required adequate post-exposure rabies post-exposure prophylaxis[12]. In future vaccination campaigns, the manual distribution of vaccines within suburban areas might be examined.

Environmental parametres

Climatic variables were derived from the WorldClim version 1.4. software [13] (University of California, Berkeley, USA). Land uses and population density were derived from the Corine Land Cover 2000 database (European Environment Agency, Copenhagen, Denmark). Boundaries of local administrative units were retrieved from the national open data catalogue (www.geodata.gov.gr; Institute for the Management of Information Systems, Athens, Greece). Distance from permanent water sources and altitude values were extracted from a digital elevation model (DEM). GIS layers were created to represent locations of towns and villages, distance to the nearest village, distance from water presence, road networks, wildlife refuges and livestock farms using the ArcGIS 10.1 GIS software (ESRI, Redlands, USA). All data layers were converted to a common projection, map extent and resolution, when used for the MaxEnt modelling. All other environmental variables were featured data type (land uses, distance from farms etc.), which were converted to raster dataset with the same resolution and cell size using the conversion tool from the spatial analyst extension. All statistical analyses were performed with SPSS 19.0 (IBM Analytics, Armonk, USA).

Environmental niche model (ENM)

In the Maxent modelling, the pixels of the study area defined the area, where distribution of the Maxent probability had been defined. Pixels with occurrence records constituted the sample points and the features were environmental parametres (climatic factors, vegetation, topography etc.). Maxent method required presence-only data, utilised both continuous and categorical data and included efficient deterministic algorithms and mathematical definitions [14]. Red fox rabies cases (n=40) and red fox negative samples of passive surveillance randomly selected (n=180) were used as occurrence points for the ENM procedure. Maximum entropy modelling, using the MaxEnt software ver. 3.3.3, (Princeton University, Princeton, USA), was employed to predict the appropriate ecological niches for red fox rabies cases [14]. Goodness of fit of the model predictions was evaluated by the mean area under the curve (AUC) of the receiver operating characteristic curve (ROC). The Jackknife procedure was used to reduce number of environmental variables to only those that showed a substantial influence on the model. According to Ceccarelli et al. [15], testing was repeated with the Jackknife procedure until all remaining variables had a positive effect on the total gain.

Descriptive statistics were extracted with frequencies and medians of the distributions. In most cases, median values were preferred instead of means, due to a non-normal distributions of the continuous variables [16]. Above data, along with coordinates of the places where rabies cases were detected in animals, were imported into the GIS environment using Arc GIS 10.1 software (ESRI, Redlands, USA). During spatial analysis, lakes, dams, rivers and urban areas were excluded (as vaccine-baits were unnecessary and should not have been dropped near human activity areas or water supply networks respectively). Environmental criteria for methodology of rabies vaccination (e.g., vegetation, distance from increased population density, water presence) were used to determine the area for implementation of the oral vaccination programme in each of the 24 regional units in Greece.

RESULTS

Cases of rabies in animals and passive surveillance data

In total, 48 animal rabies cases were diagnosed in Greece during the period October 2012 to June 2014. Details of all cases are presented in Table 1. Most cases (40/48, 83%) involved red foxes. Other animal species involved were dogs (five cases), cattle (two cases) and a cat.

Further, samples examined for passive surveillance purposes during the reference period originated from 956 animals: 174 (18%) in 2012, 551 (58%) in 2013 and 231 (24%) in 2014. Of these, most samples (488/956 [51%], 103 in 2012, 289 in 2013 and 96 in 2014) originated from red foxes. Samples from other wild animal species examined during passive surveillance originated from bat, beech marten, brown bear, European badger, European brown hare, European wild boar, golden jackal, least weasel, mink, monkey, red squirrel, various rodent species, roe deer, wildcat and wolf. Samples also originated from domestic animal species; cat, cattle, dog, goat, horse, pig and sheep. In all cases, samples examined for passive surveillance purposes from wildlife or domestic animals as above were found negative for rabies.

Of the 74 local administrative units in Greece, cases of animal rabies had been diagnosed in eight (11%). Of all cases, 80% were diagnosed in three administrative units in central northern Greece (Kilkis, Pella, Thessaloniki). Kilkis was the administrative unit, where most cases had been diagnosed, 18/48 (38%) of all cases. In total, 30 large towns and 2,893 smaller towns or villages were located within the 50 km buffer zones around each of the 48 confirmed cases of rabies. Passive surveillance samples were collected in total from 43 (58%) local administrative units throughout Greece, from 22 in 2012, 38 in 2013 and 35 in 2014, from which samples had been collected. Most samples, 428 of the 956 (45%) originated from administrative units where cases of rabies had been diagnosed. Details are presented in Table 2 and Fig. 2.

Of all cases, 75% were diagnosed in locations with an altitude of <343.5 m and within a distance of 1 km from human settlements. Median (range) altitude of locations where cases occurred, was 228 m (10-1,076) above sea level (Table 3, Fig. 3). Further, 50% of all cases were detected within a distance of 201 m from a body of water, 75% of all cases were detected within 350 m of wildlife refuges and 75% of all cases within 351 m from a provincial road and 517 m from a forest road network (Table 3).

Samples for passive surveillance were collected from locations with mean altitude of 354 m (range 0-1,492 m). In particular, samples from red foxes were collected from locations with mean altitude of 391 m (0-1,476

1
2
3 236 m), within a distance of 1,612 m (100-12600) from a human settlement; samples from dogs were collected from
4
5 237 mean altitude of 248 m (0-1,123).
6

7 238 Most cases (52%) of rabies occurred in cultivated land. Cases were also recorded in natural areas (forests,
8
9 239 shrub land, agroforestry formations) (29% of all), as well as in human settlements or surrounding areas (19%).
10
11 240 Mean (range) distance of locations where a case occurred, from a livestock farm was 201 m (0-2101). Of all
12
13 241 cases, 73% occurred in areas with increased livestock density and grazing activities according to the kernel
14
15 242 density estimator.
16
17 243

18
19
20
21 244 **Predictive ENM for red fox rabid cases**

22
23 245 The environmental variable with highest gain when used in isolation was annual precipitation, which therefore
24
25 246 appeared to have the most useful information by itself. The environmental variable that decreased gain the most
26
27 247 when it was omitted, was distance from farms, which therefore appeared to have most information that was not
28
29 248 present in other variables. Regularized training gain was 2.047, training AUC was 0.943 and unregularized
30
31 249 training gain was 2.459. Maximum entropy analysis in the region of Kilis, where most cases of the disease
32
33 250 were recorded, revealed that relative contribution of environmental variables included cultivated land in
34
35 251 combination with livestock activity. Regularised training gain was 1.536, training AUC was 0.921 and
36
37 252 unregularized training gain was 2.008.
38
39 253

40
41
42 254 **Exposure of humans to rabies**

43
44 255 The HCDCP recorded a total of 1,060 people, who were potentially exposed to rabies from October 2012 to
45
46 256 June 2014. Of these, 889 people were presented after dog bite. In most cases, exposure involved ownerless free-
47
48 257 roaming dogs (65%), whilst ownerless free-roaming cats were involved less often (8%); also, 11 (1%) people
49
50 258 had reportedly come into direct or indirect contact with vaccine baits. Details are in Table 4. Further, 721 people
51
52 259 (68% of all) had reported type III exposure, 310 (29%) type II and 19 (2%) type I exposure, whilst no
53
54 260 information regarding type of exposure was available in 10 (1%) people.
55
56
57
58
59
60

Most cases of anti-rabies treatment were administered to people in areas, where rabies animal cases had also been reported. Anti-rabies treatment administered to potentially exposed people included: vaccine administration (578 people, 54.5%), vaccine and anti-serum administration (462 people, 43.5%) or rabies anti-serum administration (20 people, 2%). In some cases, anti-serum administration was initiated while vaccine was not readily available but soon after, the implicated animal was evaluated as free of clinical signs or tested negative for rabies and prophylactic treatment was then discontinued without the addition of vaccine. In relation to year of administration, anti-rabies treatment was given to 22 (2%) people in 2012, to 574 (54%) people in 2013 and to 464 (44%) people in 2014.

Oral anti-rabies vaccination program

In 2013 and 2014, oral anti-rabies vaccination programs were implemented in 24 local administrative units, where animals had been considered to be at increased risk for infection and disease. Total surface of the area, where the oral anti-rabies vaccination was performed, was approx. 60,000 km². Average density of distributed baits, was 22 to 25 baits km⁻², depending on surface factors and year of the campaign.

DISCUSSION

Re-emergence of rabies in animals in Greece

Animal rabies has re-emerged in Greece in 2012. The outbreak of the disease has depended upon the situation in neighboring countries, all of which had reported animal rabies cases in the previous to the outbreak years [6, 18, 19]. In fact, there is a geographical similarity, in terms of altitude and land use, of the area of Kilikis (a border administrative unit of the country) with the respective unit Selemlis in FYROM, where a rabid red fox was diagnosed in 2011. Results of relevant studies have confirmed phylogenetic similarity of the Greek rabies viral strains with strains of the virus from rabies cases in FYROM, Bulgaria, Bosnia-Herzegovina, Montenegro and Serbia [7]. These findings, allied to the location of the initial cases of the disease, i.e., close to the north boundary of the country, support a hypothesis of southwards expansion of the disease from the Balkan

1
2
3 287 countries. This hypothesis could be supported by the fact that mountains can form a substantial barrier to the
4
5 288 spread of foxes into a territory, hence this could be a reason for the southwards spread of the disease, from
6
7 289 FYROM into Greece, rather than into other areas of that country. However, identification of rabies at high
8
9 290 altitudes, recently described in Bulgaria and Italy [5, 20] highlighted the permeability of natural barriers.
10
11 291 Although red foxes, the most important reservoir for maintenance of rabies in wildlife in Central or South
12
13 292 Europe [2, 21-24] are well-suited to a wide range of landscape and can potentially spread rabies in every type of
14
15 293 environment, the majority of rabid red foxes (16/40) were recorded in the area of Kilkis in a mean altitude of
16
17 294 210.1 m. Thus, it seems that landscape configuration characterized by gentle slopes influenced animal
18
19 295 movement and rabies dissemination. Rabies dissemination has occurred through fox social structures and the
20
21 296 wider dispersal of young foxes. This may be further affected by a complex interaction of various factors,
22
23 297 including environmental parameters, e.g., landscape configuration (shape of available habitat, terrain roughness,
24
25 298 disturbance etc.), which relate to dispersal of wildlife.
26
27 299 Sporadically, rabies can spill over from wildlife to domestic animals, which may then attack people. In general,
28
29 300 ownerless free-roaming dogs were responsible for the majority of cases of potential virus transmission to
30
31 301 humans (65% of cases with treatment administration). In Greece large and unrecorded numbers of ownerless
32
33 302 free-roaming dogs live in urban areas. Legal framework endorses reproductive only control schemes along with
34
35 303 proper education of owners. Municipal authorities are responsible by law for management of urban free-
36
37 304 roaming animals including the post-exposure procedure of search, capture and prompt evaluation of the
38
39 305 involved animal. High number of incidents and lack of available resources pose significant limitations in
40
41 306 performing the aforementioned tasks. Voluntary animal welfare societies collaborate with local authorities and
42
43 307 contribute to reproductive control and animal welfare issues.
44
45
46
47
48 308 Most cases of the disease have been recorded in areas of low altitude: agricultural land close to human
49
50 309 settlements. The maximum entropy analysis revealed that distance from farms contributed with the highest
51
52 310 proportion to define environmental niche profiles for rabid foxes. Notification of suspect rabies cases was,
53
54 311 according to the national rabies control and eradication program, mandatory for all parties involved. Forestry
55
56 312 officers and game wardens (who, due to the nature of their duties, mostly operate at high altitudes) were
57
58
59
60

involved in collection of dead animals and delivery to regional veterinary officers. Further, cultivated land in combination with habitats offering cover (i.e., forests, shrub land, agroforestry formations) constituted 81% of the buffer zones surrounding the locations of rabies cases. Maximum entropy analysis conducted specifically in the area of Kilgis, where most cases of rabies in foxes were recorded, revealed that cultivated land in combination with livestock activity had the highest contribution in the environmental niche profiles for rabid foxes, a finding consistent with known fox ecology [17].

Potential for transmission of rabies to people

Incidence of rabies in humans in Europe remains extremely small (less than five cases yearly). No human cases have been reported in Greece during the current outbreak in animals, as the result of applying strict preventive and effective control measures of rabies in domesticated animals, as well as in wild animal population.

Management of human exposure cases during the outbreak had some difficulties, because of (i) the large numbers of exposed people to whom treatment should have been performed, and (ii) the frequent unavailability of the animal (often ownerless free-roaming dogs or cats) involved in the incident for detailed veterinary examination. These reasons have led to increased number of treatment cases of people during the rabies outbreak. Nevertheless, cases of potential exposure after biting by domestic carnivores (dogs, cats) have also been reported, which calls for implementation of compulsory vaccinations of these animals and of proper education of owners and attack victims for post-exposure procedures (i.e. exchange of contact details, veterinary evaluation).

Certain criteria were set from Competent Health Authorities in form of algorithm to guide physicians to assess the need to administer post-exposure prophylactic anti-rabies treatment in potentially exposed humans. Due to the fact that a universal algorithm may not include every single parameter due to complexity reasons, physicians were encouraged to individualize, as possible, each case bearing in mind that irrational administration of treatments may result in unnecessary side effects and even exhaust available consumables. This could explain the fact that post-exposure prophylactic treatments were administered to humans even at areas far away and well-isolated from the locations of clinical cases of rabies.

1
2
3 339
4
5 340 **Oral anti-rabies vaccination program**
6
7 341 Oral vaccination of foxes is a proven, effective method for elimination of rabies in wildlife population. In
8
9 342 Europe, 24 countries have already implemented oral vaccination of foxes [25]. Many countries (Bosnia &
10
11 343 Herzegovina, Bulgaria, Croatia, Czech Republic, Estonia, FYROM, Germany, Kosovo, Montenegro, Poland,
12
13 344 Romania, Serbia, Slovenia) have reported significant reduction of incidence risk or elimination of rabies after
14
15 345 implementation of oral vaccination campaigns [19, 26-36]. Oral vaccination of foxes has been considered
16
17 346 successful if the number of rabies cases is reduced by over 90% in a period up to 10 years after initiation,
18
19 347 although, in many countries, this target has been achieved during a period of five years [25].
20
21
22

23 348 Extent of the outbreak indicates the need for a prevalence estimate of the virus circulation in wild animals in
24
25 349 northern Greece [8], for correct implementation of preventive measures to control further expansion of the
26
27 350 disease. Moreover, increased passive surveillance throughout the country will be necessary to detect any
28
29 351 possible new cases of the disease.
30
31

32 352
33
34 353 **Concluding remarks**
35
36 354 Application of oral vaccination program of wildlife and mandatory vaccination of domestic animals has
37
38 355 contributed to limiting the disease, with no new cases since 2014. Implementation of oral vaccination
39
40 356 campaigns of red foxes for a number of subsequent years will assure eradication of the disease among wildlife
41
42 357 and will contribute to the rabies free status of the country in the future. Rabies reoccurrence in Greece
43
44 358 emphasises the need for ongoing passive and active surveillance for important zoonotic diseases, even when
45
46 359 these are believed to have been eradicated. Enhanced passive surveillance in the cross-border areas and in areas
47
48 360 with intense human activity will permit early detection of new cases. Multidisciplinary co-operation of medical
49
50 361 and veterinary authorities and institutes is necessary for collecting all data needed for taking appropriate disease
51
52 362 control and elimination measures.
53
54

55 363
56
57 364 **FINANCIAL SUPPORT**
58
59
60

Part of the outcomes of this research that was conducted at the Faculty of Veterinary Medicine of the University of Thessaly, received partial funding from the European Union Seventh Framework Programme (2007-2013) under grant agreement no. 222633 (Wild Tech). SRUC receives support from the Scottish Government. The costs related to the passive surveillance of the disease (samples analysis) as well as the costs for the implementation of the oral rabies vaccination campaigns were co-financed by the European Commission and the Greek state budget.

CONFLICT OF INTEREST

None.

REFERENCES

1. Cliquet F, Aubert M. Elimination of terrestrial rabies in Western European countries. *Developments in Biologicals* 2004; **119**:185–204.
2. Bourhy H, *et al.* Ecology and evolution of rabies virus in Europe. *The Journal of General Virology* 1999; **80** (Pt 10):2545–2557.
3. McElhinney LM, *et al.* Molecular diversity and evolutionary history of rabies virus strains circulating in the Balkans. *The Journal of General Virology* 2011; **92**:2171–2180.
4. Picard-Meyer E, *et al.* Molecular characterisation of rabies virus strains in the Republic of Macedonia. *Archives of Virology* 2013; **158**:237–240.
5. Robardet E, *et al.* Epidemiology and molecular diversity of rabies viruses in Bulgaria. *Epidemiology and Infection* 2014; **142**:871–877.
6. Un H, *et al.* Oral vaccination of foxes against rabies in Turkey between 2008 and 2010. *Berliner Und Münchener Tierärztliche Wochenschrift* 2012; **125**:203–208.

1
2
3 389 7. **Tasioudi KE, et al.** Recurrence of animal rabies, Greece, 2012. *Emerging Infectious Diseases* 2014;
4 390 **20**:326–328.
5
6
7 391 8. **Tsiodras S, et al.** Re-emergence of animal rabies in northern Greece and subsequent human exposure,
8 392 October 2012 - March 2013. *Euro Surveillance* 2013; **18**:20474.
9
10 393 9. **Galov A, et al.** High genetic diversity and low population structure in red foxes (*Vulpes vulpes*) from
11 394 Croatia. *Mammalian Biology - Zeitschrift für Säugetierkunde* 2014; **79**:77–80.
12
13 395 10. **Birtsas P, Sokos C, Exadactylos S.** Carnivores in burned and adjacent unburned areas in a
14 396 Mediterranean ecosystem. *mammalia* 2012; **76**.
15
16 397 11. **Korou L-M, et al.** Evaluation of the first oral rabies vaccination campaign of the red foxes in Greece.
17 398 *Vaccine* 2016; **34**:41–48.
18
19 399 12. **Mähl P, et al.** Twenty year experience of the oral rabies vaccine SAG2 in wildlife: a global review.
20 400 *Veterinary Research* 2014; **45**:77.
21
22 401 13. **Hijmans RJ, et al.** Very high resolution interpolated climate surfaces for global land areas. *International*
23 402 *Journal of Climatology* 2005; **25**:1965–1978.
24
25 403 14. **Phillips SJ, Anderson RP, Schapire RE.** Maximum entropy modeling of species geographic distributions.
26 404 *Ecological Modelling* 2006; **190**:231–259.
27
28 405 15. **Ceccarelli S, et al.** Modelling the potential geographic distribution of triatomines infected by *Triatoma* virus
29 406 in the southern cone of South America. *Parasites & Vectors* 2015; **8**:153.
30
31 407 16. **Gray CD, Kinnear PR.** IBM SPSS statistics 19 made simple. Psychology Press; 2012.
32
33 408 17. **Steck F, et al.** Oral immunisation of foxes against rabies. A field study. *Zentralblatt Für*
34 409 *Veterinärmedizin. Reihe B. Journal of Veterinary Medicine. Series B* 1982; **29**:372–396.
35
36 410 18. **Kirandjiski T, et al.** First reported cases of rabies in the Republic of Macedonia. *The Veterinary Record*
37 411 2012; **170**:312.
38
39 412 19. **Ilieva, D.** Assessment of the efficiency of oral vaccination against rabies in the fox population in Bulgaria.
40 413 *Revue de Médecine Vétérinaire* 2013; **11**:521–527.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 414 20. **Mulatti P, et al.** Emergency oral rabies vaccination of foxes in Italy in 2009-2010: identification of residual
415 rabies foci at higher altitudes in the Alps. *Epidemiology and Infection* 2012; **140**:591–598.
- 416 21. **Pastoret PP, Brochier B.** Epidemiology and elimination of rabies in western Europe. *Veterinary Journal*
417 (*London, England: 1997*) 1998; **156**:83–90.
- 418 22. **Finnegan CJ, et al.** Rabies in North America and Europe. *Journal of the Royal Society of Medicine* 2002;
419 **95**:9–13.
- 420 23. **Wandeler, A.** Epidemiology and ecology of fox rabies in Europe. In: World Organisation for Animal
421 Health, eds. *Historical Perspective of Rabies in Europe and the Mediterranean Basin*. Paris, France: OIE, 2004,
422 pp. 201–214.
- 423 24. **Aikimbayev A, et al.** Fighting rabies in Eastern Europe, the Middle East and Central Asia--experts call for a
424 regional initiative for rabies elimination. *Zoonoses and Public Health* 2014; **61**:219–226.
- 425 25. **Freuling CM, et al.** The elimination of fox rabies from Europe: determinants of success and lessons for the
426 future. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 2013;
427 **368**:20120142.
- 428 26. **Smreczak M, et al.** Rabies surveillance in Poland (1992-2006). *Developments in Biologicals* 2008;
429 **131**:249–256.
- 430 27. **Hostnik P, et al.** Control of rabies in Slovenia. *Journal of Wildlife Diseases* 2006; **42**:459–465.
- 431 28. **Matouch O, et al.** Elimination of rabies in the Czech Republic. *Developments in Biologicals* 2006;
432 **125**:141–143.
- 433 29. **Cliquet F, et al.** Eliminating rabies in Estonia. *PLoS neglected tropical diseases* 2012; **6**:e1535.
- 434 30. **Müller T, et al.** Elimination of terrestrial rabies in Germany using oral vaccination of foxes. *Berliner Und*
435 *Münchener Tierärztliche Wochenschrift* 2012; **125**:178–190.
- 436 31. **European Commission.** *Report on the task force meeting of the rabies subgroup*. Zagreb, Croatia:
437 European Commission, Health and Consumers Directorate-general, Directorate G – Veterinary and International
438 Affairs, Unit G5: Veterinary programmes; 2012.

1
2
3 439 32. **European Commission.** *Programme for Surveillance, Control and Eradication of Rabies in Romania 2012.*
4
5 440 European Commission, Health and Consumers Directorate-general, Directorate G – Veterinary and International
6
7 441 Affairs, Unit G5: Veterinary programmes; 2012.
8
9
10 442 33. **Potzsch, CJ, et al.** Rabies in Montenegro: transboundary disease control from a small country’s perspective.
11
12 443 In: *Proceedings of the 13th International Symposium on Veterinary Epidemiology and Economics.* Belgium,
13
14 444 Netherlands; International Symposia on Veterinary Epidemiology and Economics, 2012, pp 232.
15
16 445 34. **Tosic K, et al.** Oral vaccination of foxes: principle and field application. *Veterinary*
17
18 446 *Journal of Republic of Srpska* 2013; **XIII**:237–244.
19
20 447 35. **Yakobson B, et al.** Implementation and monitoring of oral rabies vaccination of foxes in Kosovo
21
22 448 between 2010 and 2013—An international and intersectorial effort. *International Journal of Medical*
23
24 449 *Microbiology* 2014; **304**:902–910.
25
26
27 450 36. **Lupulovic D, et al.** First Report on the Efficiency of Oral Vaccination of Foxes against Rabies in
28
29 451 Serbia. *Zoonoses and Public Health* 2015; doi: 10.1111/zph.12196 (in press).
30
31 452
32
33 453
34
35 454
36
37 455
38
39 456
40
41 457
42
43 458
44
45 459
46
47 460
48
49 461
50
51 462
52

53 **LEGENDS OF FIGURES**
54

55 464
56
57 465 **Fig. 1.** Animal rabies cases diagnosed in Greece during the period 1951 to 1989.
58
59
60

Fig. 2. Map of geographical location of animal rabies cases diagnosed in Greece in conjunction with the negative animal samples of the passive surveillance and the rabies post-exposure prophylaxis in humans (Oct. 2012-Jun. 2014)

Fig. 3. Map of geographical location of animal rabies cases diagnosed in Greece and FYROM, in relation to topographic configuration at landscape level.

Table 1. Details, in chronological order, of confirmed cases of rabies in animals in Greece (October 2012-June 2014).

Case ID	Animal species involved	Date of occurrence	Administrative unit	Location
1	Red fox	19/10/2012	Kozani	Siatista
2	Dog	19/11/2012	Kastoria	Ieropigi
3	Red fox	07/12/2012	Kilkis	Metaksoxori
4	Red fox	12/12/2012	Kilkis	Kato Potamia
5	Dog	20/12/2012	Pella	Aridaia
6	Red fox	24/12/2012	Pella	Aridaia
7	Red fox	31/12/2012	Kilkis	Metalliko
8	Red fox	31/12/2012	Kilkis	Stavrochori
9	Red fox	31/12/2012	Pella	Loutraki
10	Red fox	14/01/2013	Kilkis	Kilkis

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

11	Red fox	18/01/2013	Kilkis	Antigonia
12	Red fox	31/01/2103	Kilkis	Koromilia
13	Red fox	20/02/2013	Kozani	Kteni
14	Cat	28/02/2013	Trikala	Agrielis
15	Red fox	28/02/2013	Kilkis	Koromilia
16	Red fox	11/03/2013	Kilkis	Vafeioxori
17	Red fox	13/03/2013	Kilkis	Xorigi
18	Red fox	22/03/2013	Trikala	Kastraki
19	Red fox	01/04/2013	Kilkis	Parochthio
20	Red fox	11/04/2013	Trikala	Platanos
21	Red fox	22/04/2013	Kilkis	Koromilia
22	Red fox	29/05/2013	Pella	Idraia
23	Red fox	06/06/2013	Thessaloniki	Sindos
24	Dog	18/06/2013	Serres	Emmanouil Pappas
25	Red fox	05/07/2013	Pella	Piperia
26	Red fox	24/07/2013	Kilkis	Lipsidrio
27	Red fox	07/08/2013	Kilkis	Kolchida
28	Red fox	28/08/2013	Pella	Mayrovouni
29	Red fox	04/10/2013	Thessaloniki	Nea Filadelfia
30	Red fox	10/10/2013	Thessaloniki	Drymos
31	Red fox	10/10/2013	Thessaloniki	Proxoma
32	Red fox	16/10/2013	Pella	Orma
33	Red fox	17/10/201	Kilkis	Pirgotos
34	Red fox	24/10/2013	Thessaloniki	Mellisochori
35	Red fox	01/11/2013	Pella	Orma
36	Cattle	20/11/2013	Kilkis	Miriophyto
37	Cattle	22/11/2013	Kilkis	Miriophyto
38	Red fox	19/12/2013	Pella	Neochori Almopias
39	Red fox	09/01/2014	Kilkis	Mayroneri
40	Red fox	15/01/2014	Thessaloniki	Modi
41	Red fox	15/01/2014	Thessaloniki	Euaggelismos Laggada
42	Red fox	15/01/2014	Thessaloniki	Neochorouda
43	Red fox	15/01/2014	Thessaloniki	Vasiloudi
44	Dog	16/01/2014	Thessaloniki	Oraiokastro
45	Dog	26/02/2014	Larissa	Elassona
46	Red fox	17/03/2014	Trikala	Agios Nikolaos
47	Red fox	31/03/2014	Trikala	Nea Pefki

48	Red fox	12/05/2014	Pella	Orma
----	---------	------------	-------	------

Table 2. Geographical locations of cases of rabies diagnosed in Greece and of animals from which samples were collected for surveillance purposes (October 2012-June 2014).

Administrative unit	Rabies cases (n)	Samples collected for surveillance purposes (n)
Kilkis	18 (38%)	64 (6.5%)
Thessaloniki	10 (21%)	102 (10.5%)
Pella	10 (21%)	40 (4%)
Trikala	5 (10%)	15 (1.5%)
Kozani	2 (4%)	81 (8.5%)
Kastoria	1 (2%)	43 (4.5%)
Larisa	1 (2%)	8 (1%)
Serres	1 (2%)	75 (8%)
Ioannina	0	65 (7%)
Drama	0	54 (5.5%)
Evros	0	54 (5.5%)
Florina	0	54 (5.5%)
Attiki	0	39 (4%)
Grevena	0	36 (4%)
Pieria	0	25 (2.5%)
Xanthi	0	24 (2.5%)

1			
2			
3	Kavala	0	23 (2.5%)
4	Halkidiki	0	22 (2.5%)
5	Karditsa	0	20 (2%)
6	Rodopi	0	19 (2%)
7	Thesprotia	0	18 (2%)
8	Imathia	0	8 (1%)
9	Preveza	0	8 (1%)
10	Evia	0	7 (0.5%)
11	Arta	0	6 (0.5%)
12	Viotia	0	6 (0.5%)
13	Lefkada	0	5 (0.5%)
14	Aetoloakarnania	0	4 (0.5%)
15	Ilia	0	4 (0.5%)
16	Lakonia	0	4 (0.5%)
17	Magnisia	0	4 (0.5%)
18	Evritania	0	3 (0.5%)
19	Fokida	0	3 (0.5%)
20	Argolida	0	2
21	Fthiotida	0	2
22	Messinia	0	2
23	Ahaia	0	1
24	Arkadia	0	1
25	Hania	0	1
26	Iraklio	0	1
27	Korinthia	0	1
28	Dodecanisa	0	1
29	Rethymno	0	1
30	Total	48 (100%)	956 (100%)
31	498		
32	499		
33	500		
34	501		
35	502		
36	503		
37	504		
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

Table 3. Environmental parameters in locations, where cases of rabies in animals had been diagnosed in Greece (October 2012-June 2014).

Parametre	Median	Range
Annual rainfall (mm)	467.50	440.00-917.00
Temperature (°C)	14.05	9.50-15.80
Distance from water (m)	660.9	0.00-2898.2
Altitude (m)	227.50	10.00-1076.00
Distance from human settlement (m)	291.42	0.00-3759.00
Distance from wildlife refuge (m)	190.00	0.00-927.07
Distance from forest road (m)	284.61	0.00-1440.00
Distance from provincial road (m)	190.00	0.00-927.00
Distance from livestock farm (m)	201.25	0-2101.00
Livestock index	0.76	0.33-2.95

1
2
3 523
4
5 524
6
7
8 525
9
10 526
11
12 527
13
14 528
15
16 529 Table 4. Details of reported source of exposure of humans, who received post-exposure prophylactic anti-rabies
17
18 530 treatment.

Source of potential exposure	Details	Number of cases	Proportion (of all cases)
Animals		1,048	99%
Dog		889	84%
	ownerless free-roaming	690	65%
	domestic	135	
	farm	38	
	hunting	11	
	not reported	15	
Cat		95	10%
	ownerless free-roaming	81	8%
	domestic	14	
Red fox		26	2%
Other animal species		38	4%
Bat		16	
Stone marten		5	
Cattle		3	
Mouse / rat		3	
Wild boar		3	
Brown bear		2	
Horse		2	
Golden jackal		1	
Pig		1	
Weasel		1	
Wolf		1	
Vaccine bait		11	1%

Direct contact	7	
Indirect contact	through dog that consumed a bait	4
Unknown		1
Total	1,060	100%

531

532

533

534

For Review Only






Figure 1: Cases of rabies diagnosed in animals in Greece during the period 1951 to 1989.
247x121mm (300 x 300 DPI)



210x296mm (300 x 300 DPI)



279x215mm (300 x 300 DPI)